

AD-A153 144

# KINETICS OF HYDROLYSIS AND PRODUCTS OF HYDROLYSIS AND PHOTOLYSIS OF TETRYL

BY ELEONORE G. KAYSER    NICHOLAS E. BURLINSON    DAVID H. ROSENBLATT  
RESEARCH AND TECHNOLOGY DEPARTMENT

22 OCTOBER 1984

Approved for public release, distribution is unlimited.

DTIC  
ELECTE  
MAY 1 1985  
S B D

DTIC FILE COPY



**NAVAL SURFACE WEAPONS CENTER**

Dahlgren, Virginia 22448 • Silver Spring, Maryland 20910

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NSWC TR 84-68	2. GOVT ACCESSION NO. AD-A153144	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) KINETICS OF HYDROLYSIS AND PRODUCTS OF HYDROLYSIS AND PHOTOLYSIS OF TETRYL		5. TYPE OF REPORT & PERIOD COVERED Final Report Feb 1980 to Dec 1981
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Eleonore G. Kayser, Nicholas E. Burlinson, and David H. Rosenblatt		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Surface Weapons Center (Code R16) White Oak Silver Spring, MD 20910		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 6.27.04A AF25
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Medical Research & Development Cmd. Attn: SGRD-RMS Fort Detrick, Frederick, MD 21701		12. REPORT DATE 22 October 1984
		13. NUMBER OF PAGES 25
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) U.S. Army Medical Bioengineering Research and Development Laboratory		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Picrate ion                      Tetryl Photolysis N-methylpicramide              Methylnitramine Tetryl                              Tetryl Hydrolysis Products Tetryl Hydrolysis                Tetryl Photolysis Products		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This study involved the determination of tetryl hydrolysis kinetics in buffered aqueous solutions at 40°C, 72°C, and 85°C, and the identification and quantification of the major products of both the light and dark hydrolysis reactions of tetryl. The kinetic data indicated a first order disappearance of tetryl for the buffered aqueous solutions at 40°C, 72°C, and 85°C, and a first order dependence on OH <sup>-</sup> ion at 40°C. The environmental hydrolysis (20°C and pH 6.8) half life of tetryl (t <sub>1/2</sub> = 302 ± 76 days) could only be		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. (Cont.)

estimated for a range of activation energies. The major components isolated from the hydrolysis study included: picrate ion, N-methylpicramide, methylnitramine, nitrate ion, and nitrite ion. Several other compounds were also formed but were not identified.

S/N 0102- LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

## FOREWORD

Since tetryl has been found as a soil contaminant associated with past Army manufacturing operations, it was necessary to determine its environmental half life and hydrolysis products to ascertain if additional clean-up procedures are necessary.

This work was funded by the U.S. Army Toxic and Hazardous Material Agency, under the direction of the U.S. Army Medical Research and Development Command (D. Rosenblatt - Project Officer, Work Unit Number 6.27.04A AF25).

This report presents data on the environmental half life of tetryl and also identifies and quantifies its major hydrolysis products.

Approved by:



J. F. PROCTOR, Head  
Energetic Materials Division

**DTIC**  
**ELECTE**  
**S** MAY 1 1985 **D**  
**B**



1/11

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

## CONTENTS

	<u>Page</u>
INTRODUCTION . . . . .	1
EXPERIMENTAL . . . . .	2
RESULTS AND DISCUSSION . . . . .	4
CONCLUSIONS AND RECOMMENDATIONS . . . . .	15

## ILLUSTRATIONS

<u>Figure</u>		<u>Page</u>
1	HPLC CHROMATOGRAMS OF AQUEOUS TETRYL IN 0.01M BORAX BUFFER, AT pH 9.0 . . . . .	5
2	HPLC CHROMATOGRAM OF STANDARD METHYLNITRAMINE, N-METHYLPICRAMIDE, TETRYL, AND PICRATE ION SOLUTION USING PIC-A REAGENT . . . . .	6
3	40°C TETRYL HYDROLYSIS RATES AT VARIOUS pH VALUES . . . . .	11
4	72°C TETRYL HYDROLYSIS RATES AT VARIOUS pH VALUES . . . . .	12
5	85°C TETRYL HYDROLYSIS RATES AT VARIOUS pH VALUES . . . . .	13

## TABLES

<u>Table</u>		<u>Page</u>
1	LC RETENTION TIMES AND RELATIVE RESPONSES FOR TETRYL AND ITS HYDROLYSIS PRODUCTS . . . . .	7
2	TETRYL HYDROLYSIS PRODUCTS . . . . .	8
3	KINETIC DATA FOR HYDROLYSIS OF TETRYL IN BUFFERED AQUEOUS SOLUTIONS . . . . .	9
4	KINETIC DATA OF TETRYL HYDROLYSIS IN BUFFERED AQUEOUS SOLUTIONS . . . . .	14

## INTRODUCTION

N,2,4,6-tetranitro-N-methylaniline (tetryl) has been used as an explosive ingredient since 1906, both in booster and main charge explosive compositions. However, in more recent systems, it is being replaced by compositions containing RDX and HMX.<sup>1</sup> Tetryl has been found as a soil contaminant associated with past Army manufacturing operations,<sup>2,3</sup> and is reported to be soluble in fresh water to the extent of 75 ppm (20°C)<sup>4</sup> and in seawater to the extent of 25 ppm (25°C).<sup>5</sup> It is also toxic with a particularly detrimental effect on the skin.<sup>6</sup>

Several researchers have reported<sup>7</sup> that tetryl hydrolyzes in mild base to picrate ion, methylamine, and nitrite ion. Methylnitramine formation is also a possibility, and due to its stability in water the latter could act as a methylating agent. Because of this stability, methylnitramine may also be a potential cause of "frame shift" mutations, although its mutagenicity has not as yet been determined. John Hoffsommer has reported on the slow hydrolysis of saturated tetryl solutions (~ 25 ppm) in seawater (pH ~ 8.1). He observed an 88 percent conversion to picric acid in 101 days at 25°C in low actinic glassware. The formation of picric acid by spectrographic measurement ( $\lambda = 352$  nm,  $\epsilon = 14,130$ )<sup>8</sup> was in agreement with the decrease noted in tetryl concentration by GC analysis.\* However, Hoffsommer suggested that biodegradation in the seawater samples was also probable.

In considering the environmental fate of tetryl in soil and groundwater, several reactions have to be considered since it is susceptible to at least three major pathways of degradation or decomposition: (1) photolysis, (2) biodegradation, and (3) hydrolysis. For this reason, the following reactions and kinetic studies were undertaken:

---

\*GC analysis of tetryl can be misleading due to thermal decomposition of tetryl in the GC injection port forming N-methylpicramide and another unidentified product.

- o A light hydrolysis\* reaction over a period of 20 days at ambient temperature.
- o Dark hydrolysis\*\* reactions for periods of 20 and 90 days at ambient temperature.
- o A borax buffered dark hydrolysis reaction over a 90-day period at ambient temperature.
- o Tetryl hydrolysis kinetics in buffered aqueous solutions (pH range 4 to 9) at 40°, 72°, and 85°C.

From the above studies, the major products of both light and dark tetryl hydrolysis reactions were identified and quantified. The kinetic data should allow the determination of a pH versus rate profile of tetryl, as well as the calculation of activation energies in order to predict hydrolysis half lives at environmental conditions (20°C and pH 6.8).

## EXPERIMENTAL

### PROCEDURE

#### Light and Dark Hydrolysis

Two solutions of tetryl were prepared in distilled water (12 mg/L). One was placed in the dark for 20 days while the other remained on the laboratory bench top in room light, both were stored at ambient temperature. The dark hydrolysis reaction was repeated with tetryl solutions (15 mg/L) in (1) 0.005 M borax buffer and (2) distilled water. The water solution was placed in the dark while the borax solution was placed in low actinic glassware on the laboratory bench top. Both solutions were again stored at ambient temperature.

#### Tetryl Hydrolysis Kinetics

The tetryl hydrolysis kinetics were determined in buffered aqueous solutions which were shielded from light. Reactions were carried out in 100 ml stoppered volumetric flasks submerged in constant-temperature baths maintained at 40°, 72°, and 85°C. Initial tetryl concentrations ranged from 1 to 20 mg/L. Two types of baths were used: (1) a Braun Thermomix 1460, 3-gallon, 1,000-watt unit, with  $\pm 0.01$  temperature stability; and (2) a modified Sargent

---

\*Light hydrolysis--a hydrolysis reaction under ambient lighting conditions.

\*\*Dark hydrolysis--a hydrolysis reaction shielded from light.

Welch Termoniter unit with 1,500 watts of proportional heating elements in a 16-gallon American Instrument Company Constant Temperature Bath with  $\pm 0.01^\circ\text{C}$  temperature stability. The NBS-calibrated thermometers were readable to  $\pm 0.02^\circ\text{C}$ . The samples were exposed to light only while cooling, prior to analysis.

#### MATERIALS, REAGENTS, AND pH DETERMINATIONS

For the majority of the kinetic runs, three NBS standard buffers were used. They were 0.01 M potassium acid phthalate (KHPth) for pH 4; equivalent moles of  $\text{KH}_2\text{PO}_4/\text{Na}_2\text{HPO}_4$  (0.005, 0.025, 0.05 M for each) for pH 6.8; and 0.01 M  $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10 \text{H}_2\text{O}$  for pH 9.18. The actual pH value was measured at each temperature ( $40^\circ$ ,  $72^\circ$ , and  $85^\circ\text{C}$ ) to allow an exact calculation of hydroxide concentration. A model number 130 Corning pH meter and a Corning Ag/AgCl combination electrode (#476115) with an accuracy of  $\pm 0.001$  pH units were used.

The pH meter was calibrated at ambient temperature with the appropriate NBS buffer standard. Twenty-five ml samples of the kinetic stock solutions were added to a small (50 ml) insulated constant temperature bath which contained the combination electrode and thermometer. The solutions, electrode, and thermometer were allowed to equilibrate for 20 minutes after the thermometer had reached temperature before a pH reading was taken. The system was sealed to prevent evaporation. Since most of the kinetic stock solutions were made up with NBS buffers to standard concentrations, values were checked against the standard pH tables. Our pH measurements were within 0.03 units of those reported.

The samples of tetryl, picric acid, and N-methylpicramide used for this study were supplied by John Hoffsommer of this Center and were determined to be 99 percent pure by LC and TLC analyses. Samples of methylnitramine were obtained from George Naufflett, also of this Center.

#### TETRYL ANALYSIS

The disappearance of tetryl was monitored via LC using a Waters ALC-440/202 instrument with a U-6K injector and an isocratic 50/50, v/v methanol/water solvent system. Flow rates were 2 ml/minute and detection was by a Waters Model 440 UV detector fixed at  $\lambda = 254 \text{ nm}$ .

A Whatman stainless steel LC column, 4.6 mm ID x 25 cm x 1/4 inch, and packed with 10  $\mu\text{m}$  C18-ODS reverse phase material was used. Tetryl concentrations were measured by comparison of integrated peak areas from the kinetic runs with calibrated external injections of known amounts of tetryl from a stock solution. The stock solution was freshly prepared for each run, kept in the dark, and refrigerated during each experiment. Twenty-five to fifty  $\mu\text{l}$  samples were removed at regular intervals from the reaction solution and injected directly into the LC, preceded and followed by the standard tetryl injection.



## PRODUCT ANALYSIS

The hydrolysis products of tetryl were identified by comparison of their LC retention times with those of authentic samples. This was verified by determining retention times on two different column support phases and in two different solvents: (1) a Whatman Reverse Phase C18 column, 4.6 mm ID x 25 cm x 1/4 inch, with 50:50 methanol/water, v/v (Figure 1), and (2) a Hewlett Packard 10  $\mu$ m RP-8, 4.6 mm x 25 cm column with 50:50 methanol/water, v/v, containing  $5 \times 10^{-3}$  M tetrabutyl ammonium phosphate and buffered at pH 7.5 (Pic-A reagent) (Figure 2). The latter column/solvent system is effective in increasing the retention time for picric acid (ion-pair liquid chromatography).<sup>9</sup> Nitrate and nitrite ions were also determined on a Dionex Ion Chromatograph Model 10, using as eluent a mixture of  $3 \times 10^{-3}$  M sodium bicarbonate and  $2.4 \times 10^{-3}$  M sodium carbonate. One milliliter samples were removed from the kinetic runs and injected directly into the IC, preceded and followed by injections of standard  $\text{NO}_2^-$  and  $\text{NO}_3^-$  solutions.

## RESULTS AND DISCUSSION

## LIGHT AND DARK HYDROLYSIS

The retention times and relative response factors of tetryl and its hydrolysis products are listed in Table 1. The results of the LC and IC analyses of the light and dark hydrolysis reaction solutions can be found in Table 2. From the data obtained in this cursory product study, it is obvious that tetryl photolyzes under ambient lighting conditions at least an order of magnitude faster than it hydrolyzes. The major detectable photoproduct is N-methylpicramide. Tetryl also reacts in concentrated sulfuric acid to form N-methylpicramide (trinitromethylaniline). This reaction proceeds fairly rapidly at elevated temperatures (e.g., 60°C and above) but slowly at ambient temperature.<sup>7</sup> The principal detectable product of the dark hydrolysis reaction in borax buffer is methylnitramine. Other identified products include: picrate ion, nitrite ion, N-methylpicramide, and nitrate ion. Several additional compounds were also formed but were not identified. The dark hydrolysis reaction in borax buffer was continued for an extra 20 days after the initial 90-day period to determine the decomposition product or products of the methylnitramine, however, no degradation of the methylnitramine was noted. Under the above photolysis conditions tetryl reacts faster than it does under environmental conditions of 40°C and a pH of 6.88 ( $t_{1/2}$  = 63 days, see Table 3).

## TETRYL KINETICS

Each tetryl hydrolysis in buffered aqueous solution followed simple pseudo first-order kinetics with respect to tetryl concentration. Several runs at 40°, 72°, and 85°C indicated good first-order log conc. versus time plots for over 300-fold changes in tetryl concentration (> 99 percent disappearance). Some runs at 40°C required over 3 months for collection of data whereas at pH 9 and

HPLC CONDITIONS  
SOLVENT: METHANOL:WATER  
(50:50, V/V)  
FLOW RATE: 2 ML/MINUTE  
UV DETECTOR: 254 NM  
COLUMN: WATERS C-18 REVERSE PHASE  
CHART SPEED: 2 CM/MINUTE




FIGURE 1. HPLC CHROMATOGRAMS OF AQUEOUS TETRYL IN 0.01M BORAX BUFFER, AT pH 9.0

HPLC CONDITIONS  
 SOLVENT: METHANOL: WATER  
 (50:50, V/V) WITH PIC-A REAGENT  
 FLOW RATE: 2 ML/MINUTE  
 UV DETECTION: 254 NM  
 HP-C<sub>8</sub> REVERSE PHASE  
 CHART SPEED: 2 CM/MINUTE




FIGURE 2. HPLC CHROMATOGRAM OF STANDARD METHYLNITRAMINE, N-METHYLPICRAMIDE, TETRYL, AND PICRATE ION SOLUTION USING PIC-A REAGENT

TABLE 1. LC RETENTION TIMES AND RELATIVE RESPONSES FOR TETRYL AND ITS HYDROLYSIS PRODUCTS

COMPOUND	RETENTION TIMES (MIN)		RELATIVE RESPONSE (WATERS)* AT 254 nm
	WATERS*	HEWLETT PACKARD**	
Tetryl	6.21	6.26	1.0
Picrate Ion	0.95	7.92	8.3
N-Methylpicramide	8.63	5.27	0.83
Methylnitramine	0.98	3.48	~0.60
NO <sub>2</sub> <sup>-</sup>	1.08	1.50	~0.002
NO <sub>3</sub> <sup>-</sup>	1.06	1.48	~0.001

\*Waters C18 Reverse Phase C18-ODS, 50:50 methanol/water, flow rate:  
2 ml/minute, UV detector: 254 nm.

\*\*Hewlett Packard 10  $\mu$ m RP-8, 50:50 methanol/water with  $5 \times 10^{-3}$  M  
tetrabutyl ammonium phosphate buffered at pH 7.5 (Pic-A reagent), flow rate:  
2 ml/minute, UV detector: 254 nm.

TABLE 2. TETRYL HYDROLYSIS PRODUCTS

HYDROLYSIS REACTION	Z TETRYL REACTED	REACTION TIME/MEDIUM	PICRATE ION	WEIGHT % OF PRODUCTS FORMED				INITIAL pH	FINAL pH
				NO <sub>2</sub> <sup>-</sup>	NO <sub>3</sub> <sup>-</sup>	METHYL- NITRAMINE	N-METHYLPICRAMIDE		
Dark	3.2	20/H <sub>2</sub> O	~2.3	← ~0.01 →		→	< 1.0	5.95	6.10
Light	95.4*	20/H <sub>2</sub> O	~3.9	9.4	35.2	0.01	41.0	5.95	4.25
Dark	5.1	90/H <sub>2</sub> O	~3.4	← 0.3 →		→	1.0	6.10	6.04
Dark	98.7*	90/0.005 M borax-H <sub>2</sub> O	28.5	4.1	3.1	66.0	4.1	8.90	8.80

\*Several other compounds were formed which were not identified

TABLE 3. KINETIC DATA FOR HYDROLYSIS OF TETRYL IN BUFFERED AQUEOUS SOLUTIONS

TEMP (°C)	pH (± 0.01)	BUFFER	NO. POINTS	$k_{obs}$ (hr <sup>-1</sup> ) x 10 <sup>-4</sup>	CORRELATION COEFFICIENT <sup>10</sup>
40	6.88	0.05 M PO <sub>4</sub>	9	4.53 ± .21	0.999
40	6.88	0.05 M PO <sub>4</sub>	8	4.93 ± .38	0.997
40	8.94	0.005 M Borax	6	446 ± 12	0.999
40	8.96	0.001 M Borax	4	388 ± 7	0.999
40	8.96	0.005 M Borax	6	437 ± 12	0.999
72	4.13	0.005 M KHPht	6	2.01 ± .14	0.999
72	4.13	0.01 M KHPht	6	2.43 ± .19	0.998
72	6.87	0.05 M PO <sub>4</sub>	6	401 ± 10	0.999
72	6.87	0.05 M PO <sub>4</sub>	7	409 ± 6	0.999
72	6.87	0.05 M PO <sub>4</sub>	7	419 ± 18	0.999
72	6.87	0.05 M PO <sub>4</sub>	7	404 ± 13	0.999
72	8.74	0.001 M Borax	5	21,300 ± 200	0.999
72	8.72	0.001 M Borax	6	20,200 ± 600	0.999
72	8.89	0.005 M Borax	3	18,800 ± 200	0.999
72	9.09	0.01 M Borax	6	19,400 ± 500	0.999
72	9.06	0.01 M Borax	3	19,600 ± 1100	0.999
85	5.67	0.025 M KH <sub>2</sub> PO <sub>4</sub>	6	261 ± 6	0.999
85	5.67	0.025 M KH <sub>2</sub> PO <sub>4</sub>	5	265 ± 8	0.999
85	6.79	0.005 M PO <sub>4</sub>	7	2840 ± 210	0.996
85	6.79	0.005 M PO <sub>4</sub>	6	2870 ± 1300	0.963
85	6.84	0.01 M PO <sub>4</sub>	6	2000 ± 130	0.998
85	7.74	0.01025 M Borax + HCl	5	3630 ± 210	0.998
85	7.77	0.01025 M Borax + HCl	5	5950 ± 3960	0.933
85	7.77	0.01025 M Borax + HCl	5	4860 ± 860	0.995
85	9.50	0.005 M Borax	6	114,000 ± 1800	0.999
85	9.54	0.005 M Borax	5	76,600 ± 19,500	0.989
85	9.61	0.01 M Borax	6	107,800 ± 4000	0.999
85	9.63	0.01 M Borax	4	87,500 ± 4300	0.999

85°C the half life of tetryl was only a few minutes. Ten-fold changes in tetryl's starting concentration (2 to 20 mg/L) caused no significant change in first-order rate constants nor was catalysis observed with any of the three buffers used. Constant ionic strength was not adhered to in this kinetic study.

The pseudo first-order rate constants ( $k_{obs}$ ) found in Table 3 were determined from plots of  $\ln$  tetryl versus time. Linear regression equations for the data were calculated by a least squares method, which also gave the errors in the pseudo first-order rate constants ( $k_{obs}$ ) and the correlation coefficients. The rate constants in Table 3 represent a 95 percent confidence interval by use of the "Student T-Test." The correlation coefficients for all but three runs were better than 0.99.

Plots of pH versus  $\log k_{obs}$  (Figures 3, 4, and 5) for the hydrolyses of tetryl at 40°, 72°, and 85°C are straight lines with slopes of 0.98, 0.83, and 0.64, respectively. Since the slope reflects the order of the reaction with respect to hydroxide, it appears that either the mechanism is changing or other reactions are occurring as the temperature increases. However, at the lower temperatures the order appears to approach 1.0. Therefore, with respect to hydrolysis at 40°C, tetryl is characterized by the standard rate expression:

$$-d \frac{[\text{tetryl}]}{dt} = k[\text{tetryl}][\text{OH}^-]$$

where in buffered solutions the following equations apply:

$$-d \frac{[\text{tetryl}]}{dt} = k_{obs}[\text{tetryl}]$$

and

$$k_{obs} = k_2[\text{OH}^-]$$

It was not possible to extract hydroxyl ion-independent constants for a possible water reaction.

Table 4 lists the observed rate constants ( $k_{obs}$ ), pH, temperature, hydroxide ion concentration, and the calculated second-order rate constants ( $k_2$ ). As can be seen, they are not constant, especially at the higher temperatures. At 40°C the average value for  $k_2 = 1754 \pm 430 \text{ M}^{-1} \text{ hr}^{-1}$ . There does not appear to be an increasing or decreasing trend in the five values at 40°C. Due to the long duration (> month) of these kinetic runs no further experiments were conducted at this temperature. Based on the available data (Figures 3, 4, and 5), the rate of tetryl hydrolysis is first order in tetryl concentration over the entire temperature range (40° to 85°C) but less than first order in hydroxide ion concentration at 72° and 85°C (see falloff in  $k_2$  values with increase in pH--Table 4). This suggests a possible competing reaction where tetryl is disappearing by a route which does not depend on hydroxide ion concentration. Evidence for a thermal first-order decomposition of tetryl is discussed below. The scope of this task did not permit further investigation of possible side reactions.




FIGURE 3. 40°C TETRYL HYDROLYSIS RATES AT VARIOUS pH VALUES






FIGURE 4. 72°C TETRYL HYDROLYSIS RATES AT VARIOUS pH VALUES




FIGURE 5. 85°C TETRYL HYDROLYSIS RATES AT VARIOUS pH VALUES

TABLE 4. KINETIC DATA OF TETRYL HYDROLYSIS IN BUFFERED AQUEOUS SOLUTIONS

TEMP (°C)	pH	$k_{obs} (hr^{-1})$	$k_2 (M^{-1}hr^{-1}) = k_{obs}/[OH^-]$	OH <sup>-</sup> CONCENTRATION (MOLES/LITER)
40	6.88	0.000453	1,805	$2.51 \times 10^{-7}$
40	6.88	0.000493	1,964	$2.51 \times 10^{-7}$
40	8.94	0.0446	1,546	$2.88 \times 10^{-5}$
40	8.96	0.0388	1,285	$3.02 \times 10^{-5}$
40	8.96	0.0437	1,447	$3.02 \times 10^{-5}$
72	4.13	0.000201	89,732	$2.24 \times 10^{-9}$
72	4.13	0.000243	108,482	$2.24 \times 10^{-9}$
72	6.87	0.0401	32,602	$1.23 \times 10^{-6}$
72	6.87	0.0409	33,252	$1.23 \times 10^{-6}$
72	6.87	0.0419	34,065	$1.23 \times 10^{-6}$
72	6.87	0.0404	32,845	$1.23 \times 10^{-6}$
72	8.72	2.02	23,192	$8.71 \times 10^{-5}$
72	8.74	2.13	23,355	$9.12 \times 10^{-5}$
72	8.89	1.88	14,687	$1.28 \times 10^{-4}$
72	9.06	1.96	10,262	$1.91 \times 10^{-4}$
72	9.09	1.94	9,509	$2.04 \times 10^{-4}$
85	5.67	0.0261	185,000	$1.41 \times 10^{-7}$
85	5.67	0.0265	188,000	$1.41 \times 10^{-7}$
85	6.79	0.284	153,000	$1.86 \times 10^{-6}$
85	6.79	0.287	154,000	$1.86 \times 10^{-6}$
85	6.84	0.200	96,000	$2.09 \times 10^{-6}$
85	7.74	0.363	22,000	$1.66 \times 10^{-5}$
85	7.77	0.595	33,000	$1.78 \times 10^{-5}$
85	7.77	0.486	27,000	$1.78 \times 10^{-5}$
85	9.50	11.40	12,000	$9.54 \times 10^{-4}$
85	9.54	7.66	7,300	$1.05 \times 10^{-3}$
85	9.61	10.78	8,800	$1.23 \times 10^{-3}$
85	9.63	8.75	6,600	$1.29 \times 10^{-3}$

Values of  $[OH^-]$  are calculated from the expression,  $\log[OH^-] = pH + \log k_w$  where  $\log k_w = -13.482$ ;  $-12.776$ ; and  $-12.523$  at  $40^\circ$ ,  $72^\circ$ , and  $85^\circ C$ , respectively.<sup>11</sup>

Attempts to extract a first-order thermal decomposition rate from the 72° and 85°C rate data were unsuccessful. This suggests a more complicated breakdown of tetryl at higher temperatures. The results of the second-order rate data (Table 4) also preclude a good Arrhenius plot, which would have allowed an accurate prediction of the hydrolysis rates under environmental conditions. In lieu of that, however, the hydrolysis rates at 40°C and pH 6.8 ( $t_{1/2} = 63$  days) give an indication of what tetryl's half life will be under environmental conditions. Assuming that the measured first-order hydrolysis rate constant ( $k_{obs}$ ) changes by a factor of two for each 10°C change in temperature, the  $t_{1/2}$  at 20°C and pH 6.8 would be 254 days. This "rule of thumb" is based on an hydrolysis activation energy ( $E_h$ ) of about 18 Kcal/mole. Since the values of  $E_h$  max range from 15 to 28 Kcal/mole, the rate constant over this range can vary by a factor of 1.8 to 3.0 for every 10°C change. Therefore, the extrapolated rate constant is only semiquantitatively correct. Based on the  $E_h$  range, the  $t_{1/2}$  range would be  $302 \pm 76$  days for tetryl hydrolysis at 20°C and a pH of 6.8.

### CONCLUSIONS AND RECOMMENDATIONS

Although the environmental hydrolysis half life of tetryl ( $t_{1/2} \approx 302 \pm 76$  days) could only be estimated for a range of activation energies, we consider this to be a realistic value for 20°C and pH 6.8. It should be kept in mind, however, that some soils<sup>12</sup> might have a strong accelerating effect on tetryl's hydrolysis rate which could drastically alter its environmental hydrolysis half life. The question of the formation of the potentially toxic methylnitramine under environmental conditions should be investigated. Methylnitramine formation was observed to be less than 5 percent in the light hydrolysis reactions but to be greater than 50 percent in the higher pH dark hydrolysis reaction. The relative hydrolysis rate of methylnitramine compared to tetryl has not been determined. Additional experiments are also needed to determine the unidentified products formed in these reactions.

## REFERENCES

1. Kaye, S. M., Ed. Encyclopedia of Explosives and Related Items, Vol. 9, U.S. Army Armament Research and Development Command, Dover, NJ, 1980, p. T-148.
2. Keirn, M. N., Stratton, C. L., Mousa, J. J., Bonds, J. D., Winegardner, D. L., Prentice, H. S., Adams, W. D., and Powell, V. J., "Environmental Survey of Alabama Army Ammunition Plant," E.S.E., Inc., Contract No. DRXTH-FS-CR-81104, Gainesville, FL, 17 Jul 1981.
3. Batzer, J., Haverl, S., Frostman, T., and Retzlaff, J., Installation Restoration Surveys: Joliet Army Ammunition Plant (JAAP), Vol. 2, Final Draft Technical Report, Report DRXTH-AS-CR-82143A, 10 Nov 1982.
4. U.S. Army Materiel Command, Engineering Design Handbook - Properties of Military Interest, Report AMC No. 706-166, Jan 1971.
5. Hoffsommer, J. C., and Rosen, J. M., "Hydrolysis of Explosives in Sea Water," Bull. Environ. Contam. Toxicol. 10 [2], 78, 1973.
6. Urbanski, T., Chemistry and Technology of Explosives, Vol. 3, (Pergamon Press, 1967), p. 56 and references cited therein.
7. Urbanski, T., Chemistry and Technology of Explosives, Vol. 3, (Pergamon Press, 1967), p. 51 and references cited therein.
8. Kamlet, M. J., Organic Electronic Spectral Data, Vol. I, (New York: Interscience Publication, 1960), p. 69.
9. Hoffsommer, J. C., Glover, D. J., and Hazzard, C. Y., "Quantitative Analysis of Polynitrophenols by Reverse-Phase Ion-Pair Chromatography," Chromatogr. 195, 1980, pp. 435-440.
10. Campbell, C., "Introduction to Simple Linear Regression," Appendix H of the 14th Monthly Progress Report, Contract DAMD 17-78-C-8073, A. D. Little, Inc.
11. Burlinson, N. E., Lee, L. A., and Rosenblatt, D. H., "Kinetics and Products of Hydrolysis of 1,2-Dibromo-3-chloropropane," Environ. Sci. Technol., 16 [9], 1982, pp. 627-632.
12. Kayser, E. G., and Burlinson, N. E., Migration of Explosives in Soil, NSWC TR 82-566, 6 Dec 1982.

## DISTRIBUTION

	<u>Copies</u>		<u>Copies</u>
Superintendent Naval Postgraduate School Monterey, CA 93940	1	Commanding Officer Naval Weapons Station Attn: Library NEDED Yorktown, VA 23690	1 1
Director Naval Research Laboratory Attn: Technical Information Section Washington, DC 20390	1	Commander Naval Sea Systems Command Attn: SEA-62A (G. Edwards) SEA-09G32 (Technical Library) SEA-64E (R. Beauregard) Washington, DC 20362	1 2 1
Commander David W. Taylor Naval Ship Research & Development Center Attn: Library Bethesda, MD 20034	1	Commander Naval Air Systems Command Washington, DC 20361	1
Commanding Officer and Director Naval Ship Research & Development Center Attn: Library Underwater Explosions Research Division Portsmouth, VA 23709	1	Chief of Naval Material Attn: Code 0323 (I. Jaffe) Washington, DC 20360	1
Commander Naval Weapons Center Attn: Technical Library Dr. A. Nielsen China Lake, CA 93555	1 1	Commanding Officer Naval Weapons Support Center Attn: Library Crane, IN 47522	1
Office of Naval Research Attn: ONR-715 Department of the Navy Arlington, VA 22217	2	Commanding Officer Naval Ammunition Depot Attn: Library McAlester, OK 74501	1
Commanding Officer Naval Weapons Evaluation Facility Attn: Library Kirtland Air Force Base Albuquerque, NM 87117	1	Commanding Officer Naval Supply Center Attn: Library Naval Biomedical Research Laboratory Oakland, CA 94625	1

## DISTRIBUTION (Cont.)

	<u>Copies</u>		<u>Copies</u>
Commanding Officer Naval Ordnance Station Attn: Library Division Indian Head, MD 20640	1	Commanding Officer Harry Diamond Laboratories 2800 Powder Mill Road Adelphi, MD 20783	1
U.S. Army Medical Bioengineering Research & Development Laboratory Attn: SGRD-UB2-IL (COL R. Carnahan)	1	Commander Eglin Air Force Base Attn: AFATL/DLDE Florida 32542	1
Jesse J. Barkley	1	Acting Director	
D. H. Rosenblatt	1	Army Material Systems	
SGRD-RMS	5	Analysis Agency	
SGRD-UBD-A	1	Attn: Library	1
UBZ-IL	1	Building 392	
SGRD-UBG	25	Aberdeen Proving Ground, MD 21105	
Fort Detrick Frederick, MD 21701		Commanding Officer U.S. Armament Research & Development Command	
Commanding General Army Material Command Attn: R&D Division	1	Attn: Library	1
Department of the Army Washington, DC 20318		Dover, NJ 07801	
Commanding General Army Missile Command Redstone Scientific Information Center		Director of the Chemical Laboratory	
Attn: Technical Library	1	Chemical Systems Laboratory	
Redstone Arsenal, AL 35809		Attn: SAREA-CL-D,	1
Commanding General Aberdeen Proving Ground		Aberdeen Proving Ground, MD 21010	
Attn: BRL	1	Commander	
Technical Library	1	U.S. Army Natick Research and Development Command	
Aberdeen, MD 21001		Attn: Library	1
Commanding Officer Picatinny Arsenal		Natick, MA 01760	
Attn: Library	1	Feltman Research Laboratory	
Dover, NJ 07801		U.S. Army Armament Research and Development Command	
Commanding Officer Edgewood Arsenal		Explosives Division	
Attn: USAEHA-OECD	1	Attn: Library	1
Aberdeen Proving Ground, MD 21010		SARPA-FR-E-C/Bldg 407	
		Dover, NJ 07801	
		Commandant	
		Academy of Health Sciences,	
		U.S. Army	
		Attn: HSHA-DCD	1
		Fort Sam Houston, TX 78234	

## DISTRIBUTION (Cont.)

	<u>Copies</u>		<u>Copies</u>
Commander U.S. Army Toxic and Hazardous Materials Agency Attn: DRXTH-TE-D Aberdeen Proving Ground, MD 21010	10	Director Los Alamos Scientific Laboratory Attn: Library GMX-7 P.O. Box 1663 Los Alamos, NM 87554	1
Defense Technical Information Center Cameron Station Alexandria, VA 22314	12	Lawrence Livermore National Laboratory University of California Attn: Technical Information Division	1
Library of Congress Attn: Gift and Exchange Division Washington, DC 20540	4	P.O. Box 808 Livermore, CA 94551	
Director U.S. Bureau of Mines Division of Explosive Technology Attn: Library 4800 Forbes Avenue Pittsburgh, PA 14213	1	The Franklin Institute Attn: Library 20th Street & Benjamin Franklin Parkway Philadelphia, PA 19103	1
U.S. Environmental Protection Agency Office of Research & Development Monitoring Techniques Division Attn: RD680 (Robert B. Medz) Washington, DC 20460	1	Jet Propulsion Laboratory California Institute of Technology Attn: Library Pasadena, CA 91109	1
NASA Langley Research Center Hampton, VA 23365	1	Stanford Research International Attn: Library Dr. Ron Spanggord 333 Ravenswood Avenue Menlo Park, CA 94025	1
NASA Scientific & Technical Information Facility P.O. Box 33 College Park, MD 20740	1	Atlantic Research Corporation Attn: Library 5390 Cherokee Avenue Alexandria, VA 22314	1
		Internal Distribution:	
		E431	9
		E432	3
		R16 (E. Kayser)	1
		R16 (N. Burlinson)	1